arXiv:1210.7588v1 [hep-ph] 29 Oct 2012

Correlations between multiplicities in rapidity and azimuthally separated windows

V.V. Vechernin

St.-Petersburg State University
E-mail: vechernin@pobox.spbu.ru

Abstract

The forward-backward multiplicity correlations in two windows separated in rapidity and azimuth are analyzed in the framework of the model with independent identical emitters (strings). Along with the long-range contribution, originating from the fluctuation in the number of emitters, the short-range contribution, originating from the correlation between multiplicities produced by a single emitter is also taken into account. The dependencies of the correlation coefficient on the rapidity and azimuthal acceptance of the windows and on the corresponding gaps between windows are studied. It is shown that the analysis of these dependencies enables to separate the contributions of two above mechanisms.

1 Introduction

In the multiparticle production in pp and AA collisions at high energies one can study the so-called forward-backward (FB) correlation between multiplicities n_F and n_B of charged particles produced in two windows ("forward" and "backward") separated by some rapidity and azimuthal gaps. We consider this correlation in the framework of the model with independent emitters [1, 2, 3], suggesting that they are strings.

In contrast with AA collisions in the case of pp collisions in the first approximation one can neglect the effects of the interaction between strings and consider them as an identical emitters at least in the central rapidity region, where each string contributes to the particle production in the whole rapidity range of the region.

In previous paper [4] we have considered only the long-range (LR) part of the correlation, originating from the fluctuation in the number of emitters. In the present note we also take into account the short-range (SR) contribution, originating from the correlation between multiplicities produced by a single emitter.

For the FB correlation coefficient b one uses the following definition:

$$b = \frac{\langle n_B n_F \rangle - \langle n_B \rangle \langle n_F \rangle}{\sqrt{D_{n_B} D_{n_F}}} \ , \tag{1}$$

which for symmetric FB windows is equivalent to

$$b = \frac{\langle n_B n_F \rangle - \langle n_B \rangle \langle n_F \rangle}{D_{n_F}} = \frac{\langle n_B n_F \rangle - \langle n_B \rangle \langle n_F \rangle}{\langle n_F^2 \rangle - \langle n_F \rangle^2} \tag{2}$$

(see also the discussion in [4]). We use the notation D_n for the variance and ω_n for the scaled variance of the n:

$$D_n = \langle n^2 \rangle - \langle n \rangle^2 \ , \qquad \omega_n = \frac{D_n}{\langle n \rangle} \ . \tag{3}$$

Similarly one can define the correlation coefficient β, which characterizes the property of the emitter - the correlation between multiplicities μ_F and μ_B, produced by a single emitter in the forward and backward windows:

$$\beta = \frac{\overline{\mu_B\mu_F} - \overline{\mu}_B\overline{\mu}_F}{\sqrt{D_{\mu_B}D_{\mu_F}}} \ , \tag{4}$$

which for symmetric FB windows is equal to

$$\beta = \frac{\overline{\mu_B\mu_F} - \overline{\mu}_B\overline{\mu}_F}{D_{\mu_F}} = \frac{\overline{\mu_B\mu_F} - \overline{\mu}_B\overline{\mu}_F}{\overline{\mu_F^2} - \overline{\mu}_F^2} \ . \tag{5}$$

2 The model

In the model with independent identical emitters [1] one assumes that $P(n_B, n_F)$ - the probability to observe simultaneously the n_F charged particles in the forward rapidity window and the n_B particles in the backward one is given by the expression:

$$P(B,F) = \sum_N w(N) \sum_{B_1,\dots,B_N} \sum_{F_1,\dots,F_N} \delta_{B\ B_1+\dots+B_N}\delta_{F\ F_1+\dots+F_N} \prod_{i=1}^{N} p(B_i,F_i) \ , \tag{6}$$

where we have used short notations:

$$F \equiv n_F \ , \qquad B \equiv n_B \ ; \qquad F_i \equiv \mu_F^i \ , \qquad B_i \equiv \mu_B^i$$

In the formula (6) the $w(N)$ is the probability to have N emitters in the given event and the $p(B_i, F_i)$ is the probability that i-th emitter produces the F_i charged particles in the forward rapidity window and the B_i particles in the backward one.

In the case of the long-range correlations considered in the previous paper [4] with a sufficiently large rapidity gap between windows one could assume that the every emitter (string) produces particles *independently* in the forward and backward windows ($p(B_i, F_i) = p_B(B_i)\ p_F(F_i)$), but in the present paper dealing with FB correlation we will not suppose this factorization. Then acting as in [4] we have

$$\langle n_B n_F\rangle \equiv \langle BF\rangle = \sum_{B,F} BF\ P(B,F) = \tag{7}$$

$$= \sum_N w(N) \sum_{B_1,\dots,B_N} \sum_{F_1,\dots,F_N} (B_1+\dots+B_N)(F_1+\dots+F_N) \prod_{i=1}^{N} p(B_i,F_i) =$$

$$= \sum_N w(N)[N\overline{\mu_B\mu_F} + (N^2-N)\overline{\mu}_B\overline{\mu}_F] = \langle N\rangle(\overline{\mu_B\mu_F} - \overline{\mu}_B\overline{\mu}_F) + \langle N^2\rangle\overline{\mu}_B\overline{\mu}_F \ ,$$

where we have used that for identical emitters at any i:

$$\sum_{B_i,F_i} B_i\ p(B_i,F_i) = \overline{\mu}_B \ , \qquad \sum_{B_i,F_i} F_i\ p(B_i,F_i) = \overline{\mu}_F \ , \qquad \sum_{B_i,F_i} B_iF_i\ p(B_i,F_i) = \overline{\mu_B\mu_F} \ .$$

By (7) the enumerator of (1) can be written as follows

$$\langle n_B n_F\rangle - \langle n_B\rangle\langle n_F\rangle = \langle N\rangle(\overline{\mu_B\mu_F} - \overline{\mu}_B\overline{\mu}_F) + D_N\overline{\mu}_B\overline{\mu}_F \tag{8}$$

and in the denominator of (1) for variances D_{n_F} and D_{n_B} we have the same expressions as in [4]:

$$D_{n_F} = \langle N\rangle D_{\mu_F} + D_N\overline{\mu}_F^2 \ , \qquad D_{n_B} = \langle N\rangle D_{\mu_B} + D_N\overline{\mu}_B^2 \ . \tag{9}$$

Below we'll consider only the case with symmetric FB windows of the same acceptance in rapidity and azimuth:

$$\Delta y_B = \Delta y_F = \Delta y \ , \qquad \Delta\varphi_B = \Delta\varphi_F = \Delta\varphi \ , \tag{10}$$

which are separated by some gap y_{gap} in rapidity and φ_{gap} in azimuth. Then the distance between the centers of this FB windows in rapidity and in azimuth is equal to

$$y = y_{gap} + \Delta y \ , \qquad \varphi = \varphi_{gap} + \Delta\varphi \ . \tag{11}$$

In this case

$$\langle n_B\rangle = \langle n_F\rangle = \langle n\rangle \ , \qquad D_{n_B} = D_{n_F} = D_n \tag{12}$$

and

$$\overline{\mu}_B = \overline{\mu}_F = \overline{\mu} \ , \qquad D_{\mu_B} = D_{\mu_F} = D_\mu \ . \tag{13}$$

For symmetric windows the formula (8) leads to the following expression for the FB correlation coefficient b:

$$b = \frac{\beta + a}{1 + a} \ , \tag{14}$$

where β is given by (4) and a is equal to

$$a = \frac{\omega_N}{\omega_\mu}\overline{\mu} \ , \qquad \omega_N = \frac{D_N}{\langle N\rangle} \ , \qquad \omega_\mu = \frac{D_\mu}{\overline{\mu}} \ . \tag{15}$$

Recall that $\langle N\rangle$ is the mean number of emitters (strings), the $\overline{\mu}$ is the mean number of charged particles produced in the given acceptance $\Delta y\Delta\varphi$ by one emitter, D_N and D_μ are the corresponding variances. In the central rapidity region one can also consider that

$$\overline{\mu} = \mu_0\Delta y\Delta\varphi/2\pi \ , \tag{16}$$

where μ_0 is the average rapidity density of the charged particles produced by one string in this region.

Clear that in the case $\beta = 0$, i.e. without correlation between multiplicities μ_F and μ_B produced by a single emitter in the forward and backward windows (4), we come back to our formula

$$b^{LR} = \frac{a}{1 + a} \ , \tag{17}$$

obtained in [2, 3, 4] for the long-range correlation. So we can rewrite (14) as

$$b = b^{SR} + b^{LR} \ , \tag{18}$$

where

$$b^{SR} = \frac{\beta}{1 + a} \ . \tag{19}$$

3 Long-range correlation for poissonian emitters

Usually one supposes [1, 3, 5] the poissonian distribution of charge particles produced by one string in the given azimuth and rapidity acceptance $\Delta y \Delta\varphi$. In this case $\omega_\mu = 1$ and by (15)

$$a = \omega_N \overline{\mu} = \omega_N \mu_0 \Delta y \Delta\varphi / 2\pi = \alpha \Delta y \Delta\varphi / 2\pi \ , \tag{20}$$

with

$$\alpha = \omega_N \mu_0 \ . \tag{21}$$

Note that the fluctuation in the number of strings in pp and especially in AA collisions [6] is not poissonian and hence $\omega_N \neq 1$. Its value depends on the collision energy. Recall also that μ_0 is the average rapidity density of the charged particles produced by one string.

By (17) and (20) we see that the strength of long-range correlation b^{LR}, originating from the fluctuation in the number of emitters, depends only on this one parameter α and the acceptance $\Delta y \Delta\varphi$ of the windows:

$$b^{LR} = \frac{\alpha \Delta y \Delta\varphi / 2\pi}{1 + \alpha \Delta y \Delta\varphi / 2\pi} \ . \tag{22}$$

4 Short-range correlation

Contrary to the LR contribution (22) the strength of the short-range correlation b^{SR} (19) depends also on the property of the emitter, namely it is proportional to the correlation strength β between multiplicities μ_F and μ_B, produced by a single emitter in the forward and backward windows (5). The strength β of the correlation depends on the details of the string fragmentation into hadrons.

In Schwinger mechanism the string breaks into the pieces by the creation of the quark-antiquark pairs. In very first approximation one can consider that the string decomposes into the segments of the same length y_0 in rapidity. This assumption is consistent with the fact of the homogenous rapidity distribution of produced charged particles. Each i-th string segment has some transverse momentum $\mathbf{p}_i$, which is equal to the sum of the transverse momenta of the quark and antiquark at its ends

$$\mathbf{p}_i = \mathbf{p}_i^q + \mathbf{p}_i^{\overline{q}} \ . \tag{23}$$

Further fragmentation of the string segment of the length y_0 with the transverse momentum $\mathbf{p}_i$ into hadrons goes through the formation of the minijet and the decay of resonances. If the particles from the fragmentation of the same minijet (same string segment) are registered in both our windows (forward and backward), then we'll have the additional short-range near side correlation with some characteristic length in rapidity y_1 and width φ_1 in azimuth.

The additional away side correlation will also take place. It arises because of the transverse momentum conservation in the process of the quark-antiquark pair creation. Due to this reason we have

$$\mathbf{p}_i^q = -\mathbf{p}_{i-1}^{\overline{q}} \ , \qquad \mathbf{p}_i^{\overline{q}} = -\mathbf{p}_{i+1}^q \ . \tag{24}$$

But because we have to add some arbitrary momentum $\mathbf{p}_{i-1}^q$ to the momentum $\mathbf{p}_{i-1}^{\overline{q}}$ to obtain the transverse momentum of the *(i–1)*-th string segment and some arbitrary momentum $\mathbf{p}_{i+1}^{\overline{q}}$ to the momentum $\mathbf{p}_{i+1}^q$ to obtain the transverse momentum of the *(i+1)*-th string segment

$$\mathbf{p}_{i-1} = \mathbf{p}_{i-1}^q + \mathbf{p}_{i-1}^{\overline{q}} \ , \qquad \mathbf{p}_{i+1} = \mathbf{p}_{i+1}^q + \mathbf{p}_{i+1}^{\overline{q}} \ , \tag{25}$$



Figure 1: The forward-backward (FB) correlation coefficient b (2), with taking into account the long-range (LR) and short-range (SR) contributions, calculated by formulae (18), (19), (22) and (26), for the windows acceptance $\Delta y = 0.2$ and $\Delta\varphi = \pi/4$.

we expect in the away side the two broader (than in the near side) correlation peaks displaced on the $\pm y_0$ in the rapidity with the characteristic length $y_2 > y_1$ in rapidity and the width $\varphi_2 > \varphi_1$ in azimuth.

Such behavior of the β - the correlation coefficient between multiplicities μ_F and μ_B, produced by a single string in the forward and backward windows (5) - can be modeled by the following function,

$$\beta = \beta(y,\varphi) = a_1 e^{-\frac{|y|}{y_1}} e^{-\frac{\varphi^2}{\varphi_1^2}} + a_2 \left(e^{-\frac{|y-y_0|}{y_2}} + e^{-\frac{|y+y_0|}{y_2}} \right) e^{-\frac{(\pi-\varphi)^2}{\varphi_2^2}} \quad , \tag{26}$$

which depends on the distances y and φ between the centers of our FB windows in rapidity and in azimuth (11). We imply that $\varphi \in [0,\pi]$. If $\varphi_{gap} + \Delta\varphi > \pi$, then we consider that $\varphi = 2\pi - (\varphi_{gap} + \Delta\varphi)$.

5 Behavior of the resulting FB correlation strength

An example of the resulting behavior of the FB correlation coefficient b (2) as a function of the distance y between the backward and forward windows in rapidity with taking into account the LR and SR contributions, calculated by formulae (18), (19), (22) and (26) at the different values of the distance φ between the windows in azimuth, is presented in Fig.1 for the FB windows acceptance $\Delta y = 0.2$ and $\Delta\varphi = \pi/4$ at the following values of the parameters:

$\alpha = 2.1$, $y_0 = 0.9$,
$a_1 = 0.06$, $y_1 = 0.55$, $\varphi_1 = 1.0$,
$a_2 = 0.01$, $y_2 = 2.0$, $\varphi_2 = 2.0$.
In Fig.1 we see that even at the rather small value of the acceptance $\Delta y \Delta\varphi/2\pi = 0.2/8 = 0.025$ the contribution of the LR correlation mechanism is the dominating one.

6 Conclusion

By the formula (14) we see that there are two contributions to the forward-backward (FB) multiplicity correlation coefficient b (2).

The long-range (LR) contribution (17) originates from the fluctuation in the number of emitters. This contribution depends only on the one parameter, which by (20) is proportional to the scaled event-by-event variance of the number of emitters and the mean number of charged particles produced in the window by one emitter. This results in strong dependence of the LR contribution on the window acceptance $\Delta y \Delta\varphi$ (22).

The short-range (SR) contribution (19) originate from the correlation between multiplicities produced by a single emitter (5). Its strength β (26) as discussed in Sec.4 depends on the distances y and φ between the centers of the backward and forward windows in rapidity and in azimuth (11).

So we conclude that the analysis of the dependencies of the forward-backward (FB) multiplicity correlation coefficient b (2) on the acceptance $\Delta y \Delta\varphi$ and the distances y and φ between the backward and forward windows in rapidity and in azimuth enables on the base of the relation (14) to separate the contributions of the SR and LR mechanisms.

The author thanks M.A. Braun, G.A. Feofilov and I. Altsybeev for useful discussions. The work was supported by the RFFI grant 12-02-00356-a.

References

[1] M.A. Braun, C. Pajares and V.V. Vechernin, Phys. Lett. **B493**, 54 (2000).

[2] V.V. Vechernin, R.S. Kolevatov, hep-ph/0304295 (2003);
Vestnik SPbU, ser.4, no.2, 12 (2004).

[3] M.A. Braun, R.S. Kolevatov, C. Pajares, V.V. Vechernin, Eur. Phys. J. **C32**, 535 (2004).

[4] V.V. Vechernin. Proceedings of the Baldin ISHEPP XX, vol.2, JINR, Dubna, 2011, 10-15;
arXiv:1012.0214 [hep-ph] (2010).

[5] V.V. Vechernin, R.S. Kolevatov, Physics of Atomic Nuclei **70**, 1797 (2007).

[6] V. V. Vechernin and H. S. Nguyen, Phys. Rev. C **84**, 054909 (2011).